

Note on the Controllability of Molecular Systems

Viswanath Ramakrishna

Department of Mathematical Sciences and Center for Signals, Systems and

Telecommunications

University of Texas at Dallas

Richardson, TX 75083 USA

email: vish@utdallas.edu

Supported in part by NSF - DMS-0072415

ABSTRACT: The purpose of this brief note is to show that the set of states reachable from a given initial state for a finite dimensional quantum system is equal to the orbit under the Lie group generated by the Lie algebra generated by the internal and external Hamiltonians, *even if this Lie group is not compact*. This result has been explicitly mentioned in several talks by the author. This note is not being submitted for publication in any journal.

1 A Slight Improvement

Consider a finite- dimensional quantum system:

$$\dot{c} = Ac + Bc\epsilon(t), c(0) = c_0 \quad (1.1)$$

Here c evolves on the complex sphere in some C^N , and A and B are two skew-Hermitian matrices. It is well known that the set of states reachable from c_0 , via the application of admissible controls, equals the orbit through c_0 of the group, G corresponding to the Lie algebra generated by the matrices A and B , *provided this Lie group is compact*. See [5, 6, 4] for instance. *This result is valid even if G is not compact*. This result has been mentioned explicitly by the author in several talks and grant proposals, though the result has been recorded only cryptically in [7, 8] (primarily because it offers no significant practical improvements). Since such a nominal improvement may, nevertheless, be of interest, a discussion of it is provided below.

To that end, notice that A may be assumed to be diagonal, $A = \text{diag}(i\lambda_1, i\lambda_2, \dots, i\lambda_N)$. Writing $c_k = a_k + ib_k, k = 1, \dots, N$ and viewing the given quantum system as evolving on R^{2N} , it follows that the drift portion of the system, i.e., the vector field Ac , is a classical Hamiltonian system with Hamiltonian, $H = \sum_{k=1}^N \lambda_k(a_k^2 + b_k^2)$ (under the standard symplectic structure of R^{2N}).

Now the quantum system evolves on the sphere $S = \{\sum_{i=1}^N |c_i|^2 = 1\}$. S is of course also the set $\{\sum_{i=1}^N a_i^2 + b_i^2 = 1\}$. Thus S is a bounded set and further the orbit of G through the initial condition in S is also bounded, even if it is not compact. Thus, the quantum system can be viewed as a nonlinear control system on a bounded manifold, whose drift is a classical Hamiltonian system. Since this implies that the drift is weakly positively Poisson stable, the desired conclusion follows from [2](see also [1]). Indeed, these articles show that, under the weak positively Poisson stability of the drift, the reachable set from the initial condition is exactly the leaf of the foliation associated to the drift and control vector fields. But this leaf is nothing but the orbit of G through the initial condition. It also implies that the reachable set of the associated unitary generator system from the identity matrix is G (cf., [3]), since if this is not the case, the reachable set of the quantum system could not be the orbit of G through the initial condition.

In the opinion of the author, this improvement is of mathematical interest only. The result is of relevance in the situation that the quantum system is not completely controllable, i.e., when the orbit of G is not all of S . However, for the result to have practical value it is desirable to describe the orbit explicitly. One way to do this is to use invariant theory and describe the orbit as a level set. However, invariant theory is tractable only for compact and reductive groups. The author is not aware of any practical situation in which G is not compact but is reductive.

References

- [1] W. P. Dayawansa, unpublished lecture notes, University of Maryland, 1994.

- [2] K. Y. Lian, L. S. Wang and L. C. Fu, IEEE Trans on Automatic Control, *39*, 2426, 1994.
- [3] F. Albertini and D. D'Alessandro, arXiv, quant-ph, 0106128, 2001.
- [4] S. G. Schirmer and J. V. Leahy, arXiv, quant-ph 0010032, 2000.
- [5] V. Jurdjevic and H. Sussmann, Journal of Diff Equations, *12*, 313, 1972.
- [6] V. Ramakrishna et al., *Physical Review A*, Vol 51, No 2, 960, 1995.
- [7] V. Ramakrishna and H. Rabitz ‘Control of Molecular Dynamics”, in **Systems Modelling and Optimization**, M.Polis, A. Dontchev, P.Kall, I.Lasiecka and A. Olbrot, Eds., Chapman & Hall/CRC Research Notes in Mathematics, Vol. 396, pp.495-503, 1999.
- [8] V. Ramakrishna and H. Rabitz, ‘Control Applications in Chemical Physics”, V. Ramakrishna and H. Rabitz, Proceedings of II Mexican School on Gravitation and Mathematical Physics, December 6th 1996, Tlaxcala, Mexico, 35th Annivesary of Cinvestav (proceedings will be available only electronically, for details of the site where they will be posted please contact Professor Tonatiuh Matos, tmatos@fis.cinvestav.mx).